

# Modified Equations for Variational Integrators

Mats Vermeeren

Technische Universität Berlin

IIMAS, UNAM

August 23, 2018



Discretization in  
Geometry and Dynamics  
SFB Transregio 109



Berlin  
Mathematical  
School

# Table Of Contents

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]

- [1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.
- [2] V. [Modified equations for variational integrators](#). Numer. Math. 137:1001, 2017.
- [3] V. [Numerical precession in variational discretizations of the Kepler problem](#).  
[arXiv:1602.01049](#).
- [4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). [arXiv:1709.09567](#).

# Table Of Contents

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]

- [1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.
- [2] V. [Modified equations for variational integrators](#). Numer. Math. 137:1001, 2017.
- [3] V. [Numerical precession in variational discretizations of the Kepler problem](#). arXiv:1602.01049.
- [4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). arXiv:1709.09567.

# Continuous Lagrangian Mechanics

Lagrange function:  $L : \mathbb{R}^{2N} \cong TQ \rightarrow \mathbb{R} : (q, \dot{q}) \mapsto L(q, \dot{q})$ .

Solutions are curves  $q(t)$  that minimize (or are critical points of) the [action](#)

$$S = \int_{t_0}^{t_1} L(q(t), \dot{q}(t)) dt$$

where the integration interval  $[t_0, t_1]$  and the boundary values  $q(t_0)$  and  $q(t_1)$  are fixed.

$$\begin{aligned} 0 = \delta S &= \int_{t_0}^{t_1} \frac{\partial L}{\partial q} \delta q + \frac{\partial L}{\partial \dot{q}} \delta \dot{q} dt \\ &= \int_{t_0}^{t_1} \left( \frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} \right) \delta q dt + \left( \frac{\partial L}{\partial \dot{q}} \delta q \right) \Big|_{t_0}^{t_1} \end{aligned}$$

Euler-Lagrange Equation:  $\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} = 0$ .

# Continuous Lagrangian Mechanics

Lagrange function:  $L : \mathbb{R}^{2N} \cong TQ \rightarrow \mathbb{R} : (q, \dot{q}) \mapsto L(q, \dot{q})$ .

Solutions are curves  $q(t)$  that minimize (or are critical points of) the [action](#)

$$S = \int_{t_0}^{t_1} L(q(t), \dot{q}(t)) dt$$

where the integration interval  $[t_0, t_1]$  and the boundary values  $q(t_0)$  and  $q(t_1)$  are fixed.

$$\begin{aligned} 0 = \delta S &= \int_{t_0}^{t_1} \frac{\partial L}{\partial q} \delta q + \frac{\partial L}{\partial \dot{q}} \delta \dot{q} dt \\ &= \int_{t_0}^{t_1} \left( \frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} \right) \delta q dt + \left( \frac{\partial L}{\partial \dot{q}} \delta q \right) \Big|_{t_0}^{t_1} \end{aligned}$$

Euler-Lagrange Equation:  $\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} = 0$ .

# Legendre transformation

Relates Hamiltonian and Lagrangian formalism:

$$p\dot{q} = H(q, p) + L(q, \dot{q}).$$

Differentiating w.r.t.  $\dot{q}$ ,  $p$  and  $q$ ,

$$p = \frac{\partial L}{\partial \dot{q}}$$

$$\dot{q} = \frac{\partial H}{\partial p}$$

$$0 = \frac{\partial H}{\partial q} + \frac{\partial L}{\partial q} = \left( \frac{\partial H}{\partial q} + \dot{p} \right) + \left( \frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} \right),$$

establishes equivalence between Lagrangian and **Hamiltonian equations of motion**.

# Legendre transformation

Relates Hamiltonian and Lagrangian formalism:

$$p\dot{q} = H(q, p) + L(q, \dot{q}).$$

Differentiating w.r.t.  $\dot{q}$ ,  $p$  and  $q$ ,

$$p = \frac{\partial L}{\partial \dot{q}}$$

$$\dot{q} = \frac{\partial H}{\partial p}$$

$$0 = \frac{\partial H}{\partial q} + \frac{\partial L}{\partial q} = \left( \frac{\partial H}{\partial q} + \dot{p} \right) + \left( \frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} \right),$$

establishes equivalence between Lagrangian and **Hamiltonian equations of motion**.

Only works if the Lagrangian is nondegenerate:  $\left| \frac{\partial^2 L}{\partial \dot{q}^2} \right| \neq 0$

Hamiltonian systems preserve the symplectic 2-form  $\omega = \sum_i dp_i \wedge dq_i$ .

# Symplectic structure

Let  $\Phi_t$  be the flow of a Hamiltonian system, i.e.

$$\Phi_0(q, p) = (q, p)$$

and

$$\frac{d}{dt}\Phi_t(q, p) = \left( \frac{\partial H}{\partial p}(\Phi_t(q, p)), -\frac{\partial H}{\partial q}(\Phi_t(q, p)) \right).$$

Then for each  $t$ ,  $\Phi_t$  is a symplectic map,

$$\Phi_t^* \omega = \omega,$$

where  $\omega$  is the canonical symplectic form

$$\omega = \sum_i dq_i \wedge dp_i.$$

## Definition

A *symplectic integrator* is a discretization (in time) of a Hamiltonian systems, such that each discrete time-step is given by a symplectic map.



# Discrete Lagrangian mechanics

Lagrange function:  $L : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R} : (x, \tilde{x}) \mapsto L(x, \tilde{x})$ .

Solutions are **discrete curves**  $x = (x_0, x_1, \dots, x_n)$  that are critical points of the **action**

$$S_{\text{disc}} = \sum_{j=1}^n h L_{\text{disc}}(x_{j-1}, x_j)$$

Euler-Lagrange equation:

$$D_2 L_{\text{disc}}(x_{j-1}, x_j) + D_1 L_{\text{disc}}(x_j, x_{j+1}) = 0,$$

where  $D_1, D_2$  denote the partial derivatives of  $L_{\text{disc}}$ .

## Definition

A **variational integrator** for a continuous system with Lagrangian  $\mathcal{L}$  is a discrete Lagrangian system with

$$L_{\text{disc}}(x(t-h), x(t)) \approx \mathcal{L}(x(t), \dot{x}(t)),$$

and hence  $S_{\text{disc}} \approx S$ .

# Equivalence

## Theorem

*If the Lagrangian/Hamiltonian is regular, variational and symplectic integrators are equivalent.*

**Proof.** the discrete Lagrangian is a generation function of the symplectic map describing one time step,

$$\begin{aligned}p_j &= -hD_1 L_{\text{disc}}(x_j, x_{j+1}) \\ p_{j+1} &= hD_2 L_{\text{disc}}(x_j, x_{j+1})\end{aligned}$$



## Example: Störmer-Verlet method

Consider a mechanical system  $\ddot{x} = -U'(x)$  with Lagrangian

$$\mathcal{L}(x, \dot{x}) = \frac{1}{2} \langle \dot{x}, \dot{x} \rangle - U(x)$$

The Störmer-Verlet discretization is given by the discrete Lagrangian

$$L_{\text{disc}}(x_j, x_{j+1}) = \frac{1}{2} \left\langle \frac{x_{j+1} - x_j}{h}, \frac{x_{j+1} - x_j}{h} \right\rangle - \frac{1}{2} U(x_j) - \frac{1}{2} U(x_{j+1})$$

Its discrete Euler-Lagrange equation is  $\frac{x_{j+1} - 2x_j + x_{j-1}}{h^2} = -U'(x_j)$

## Example: Störmer-Verlet method

Consider a mechanical system  $\ddot{x} = -U'(x)$  with Lagrangian

$$\mathcal{L}(x, \dot{x}) = \frac{1}{2} \langle \dot{x}, \dot{x} \rangle - U(x)$$

The Störmer-Verlet discretization is given by the discrete Lagrangian

$$L_{\text{disc}}(x_j, x_{j+1}) = \frac{1}{2} \left\langle \frac{x_{j+1} - x_j}{h}, \frac{x_{j+1} - x_j}{h} \right\rangle - \frac{1}{2} U(x_j) - \frac{1}{2} U(x_{j+1})$$

Its discrete Euler-Lagrange equation is  $\frac{x_{j+1} - 2x_j + x_{j-1}}{h^2} = -U'(x_j)$

Abstract notation:  $\psi(x_{j-1}, x_j, x_{j+1}; h) = 0$  with

$$\psi(x_{j-1}, x_j, x_{j+1}; h) = \frac{x_{j+1} - 2x_j + x_{j-1}}{h^2} + U'(x_j)$$

Symplectic equivalent:

$$x_{j+1} = x_j + hp_j - \frac{h^2}{2} U'(x_j)$$

$$p_{j+1} = p_j - \frac{h}{2} U'(x_j) - \frac{h}{2} U'(x_{j+1})$$


# Table Of Contents

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]


- [1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.
- [2] V. [Modified equations for variational integrators](#). Numer. Math. 137:1001, 2017.
- [3] V. [Numerical precession in variational discretizations of the Kepler problem](#). arXiv:1602.01049.
- [4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). arXiv:1709.09567.

# Modified Equations

Exact solution of a differential equation:



Numerical solution with a variational integrator:




Notice conservation of Energy:

- ▶ Easy to prove for (continuous) Hamiltonian systems
- ▶ Follows by Noether's theorem from invariance under time-translation of the Lagrangian
- ▶ Symplectic/variational integrators very nearly preserve energy. Why?


# Modified Equations

## Conservation of Energy:

- ▶ Easy to prove for (continuous) Hamiltonian systems
- ▶ Follows by Noether's theorem from invariance under time-translation of the Lagrangian
- ▶ Symplectic/variational integrators very nearly preserve energy. Why?



Idea of proof: find a **modified equation**, a differential equation with solutions that interpolate the numerical solutions:



# Modified Equations

Modified equations are usually given by power series.  
Often they do not converge.

## Definition

*The differential equation  $\ddot{x} = f(x, \dot{x}; h)$ , where*

$$f(x, \dot{x}; h) \simeq f_0(x, \dot{x}) + hf_1(x, \dot{x}) + h^2 f_2(x, \dot{x}) + \dots$$

*is a **modified equation** for the second order difference equation  $\Psi(x_{j-1}, x_j, x_{j+1}; h) = 0$  if, for every  $k$ , every solution of the truncated differential equation*

$$\begin{aligned}\ddot{x} &= \mathcal{T}_k(f_h(x, \dot{x})) \\ &= f_0(x, \dot{x}) + hf_1(x, \dot{x}) + h^2 f_2(x, \dot{x}) + \dots + h^k f_k(x, \dot{x}).\end{aligned}$$

*satisfies*

$$\Psi(x(t-h), x(t), x(t+h); h) = \mathcal{O}(h^{k+1}).$$



# Modified Equations for symplectic integrators

Symplectic integrators are known to very nearly preserve energy, because

## Theorem

The modified equation for a symplectic integrator is a Hamiltonian equation.

Can we arrive at a similar result purely on the Lagrangian side?

Are modified equations for variational integrators Lagrangian?

# Table Of Contents

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]

- [1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.
- [2] V. [Modified equations for variational integrators](#). Numer. Math. 137:1001, 2017.
- [3] V. [Numerical precession in variational discretizations of the Kepler problem](#). arXiv:1602.01049.
- [4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). arXiv:1709.09567.

## General idea

Look for a modified Lagrangian  $\mathcal{L}_{\text{mod}}(x, \dot{x})$  such that the discrete Lagrangian  $L_{\text{disc}}$  is its **exact discrete Lagrangian**, i.e.

$$\int_{(j-1)h}^{jh} \mathcal{L}_{\text{mod}}(x(t), \dot{x}(t)) dt = hL_{\text{disc}}(x((j-1)h), x(jh)).$$

The Euler-Lagrange equation of  $\mathcal{L}_{\text{mod}}$  will then be the modified equation.

The best we can hope for is to find such a modified Lagrangian up to an error of arbitrarily high order in  $h$ .

# The discrete Lagrangian evaluated on a continuous curve

We can write the discrete Lagrangian as a function of  $x$  and its derivatives, all evaluated at the point  $jh - \frac{h}{2}$ ,

$$\begin{aligned}\mathcal{L}_{\text{disc}}[x] &= L_{\text{disc}}\left(x - \frac{h}{2}\dot{x} + \frac{1}{2}\left(\frac{h}{2}\right)^2\ddot{x} - \dots, \right. \\ &\quad \left. x + \frac{h}{2}\dot{x} + \frac{1}{2}\left(\frac{h}{2}\right)^2\ddot{x} + \dots, \quad h\right). \\ &= L_{\text{disc}}(x_{j-1}, x_j; h)\end{aligned}$$

Here and in the following:

- ▶  $[x]$  denotes dependence on  $x$  and any number of its derivatives,
- ▶ we evaluate at  $t = jh - \frac{h}{2}$  whenever we omit the variable  $t$ , i.e.  $x = x(jh - \frac{h}{2})$ ,
- ▶  $x_j = x(jh)$  and  $x_{j-1} = x((j-1)h)$ .

# A truly continuous Lagrangian

We want to write the discrete action

$$S_{\text{disc}} = \sum_{j=1}^n h L_{\text{disc}}(x_{j-1}, x_j) = \sum_{j=1}^n h \mathcal{L}_{\text{disc}} \left[ x \left( jh - \frac{h}{2} \right) \right]$$

as an integral.

## Lemma (Euler-MacLaurin formula)

For any smooth function  $f : \mathbb{R} \rightarrow \mathbb{R}^N$  we have

$$\begin{aligned} \sum_{j=1}^n h f \left( jh - \frac{h}{2} \right) &\simeq \int_0^{nh} \sum_{i=0}^{\infty} h^{2i} (2^{1-2i} - 1) \frac{B_{2i}}{(2i)!} f^{(2i)}(t) dt \\ &= \int_0^{nh} \left( f(t) - \frac{h^2}{24} \ddot{f}(t) + \frac{7h^4}{5760} f^{(4)}(t) + \dots \right) dt, \end{aligned}$$

where  $B_i$  are the Bernoulli numbers.

The symbol  $\simeq$  indicates that this is an asymptotic series.

# A truly continuous Lagrangian

## Definition

We call

$$\begin{aligned}\mathcal{L}_{\text{mesh}}[x(t)] &= \mathcal{L}_{\text{disc}}[x(t)] + \sum_{i=1}^{\infty} (2^{1-2i} - 1) \frac{h^{2i} B_{2i}}{(2i)!} \frac{d^{2i}}{dt^{2i}} \mathcal{L}_{\text{disc}}[x(t)] \\ &= \mathcal{L}_{\text{disc}}[x(t)] - \frac{h^2}{24} \frac{d^2}{dt^2} \mathcal{L}_{\text{disc}}[x(t)] + \frac{7h^4}{5760} \frac{d^4}{dt^4} \mathcal{L}_{\text{disc}}[x(t)] + \dots\end{aligned}$$

the *meshed modified Lagrangian* of  $L_{\text{disc}}$ .

Formally, the meshed modified Lagrangian satisfies

$$\int \mathcal{L}_{\text{mesh}}[x(t)] dt = \sum h L_{\text{disc}}(x_j, x_{j+1})$$

where  $x_j = x(jh)$ .

# A truly continuous Lagrangian

## Definition

We call

$$\begin{aligned}\mathcal{L}_{\text{mesh}}[x(t)] &= \mathcal{L}_{\text{disc}}[x(t)] + \sum_{i=1}^{\infty} (2^{1-2i} - 1) \frac{h^{2i} B_{2i}}{(2i)!} \frac{d^{2i}}{dt^{2i}} \mathcal{L}_{\text{disc}}[x(t)] \\ &= \mathcal{L}_{\text{disc}}[x(t)] - \frac{h^2}{24} \frac{d^2}{dt^2} \mathcal{L}_{\text{disc}}[x(t)] + \frac{7h^4}{5760} \frac{d^4}{dt^4} \mathcal{L}_{\text{disc}}[x(t)] + \dots\end{aligned}$$

the *meshed modified Lagrangian* of  $L_{\text{disc}}$ .

Formally, the meshed modified Lagrangian satisfies

$$\int \mathcal{L}_{\text{mesh}}[x(t)] dt = \sum h L_{\text{disc}}(x_j, x_{j+1})$$

where  $x_j = x(jh)$ .

Are we finished?

$\mathcal{L}_{\text{mesh}}[x]$  depends on many more derivatives than the original  $\mathcal{L}(x, \dot{x})$ .

# The meshed variational problem


## Definition

*classical variational problem*: find critical curves of some action  $\int_a^b \mathcal{L}[x(t)] dt$  in the set of smooth curves  $\mathcal{C}^\infty$ .


*meshed variational problem*: find critical curves of some action  $\int_a^b \mathcal{L}[x(t)] dt$  in the set of piecewise smooth curves that are consistent with a mesh of size  $h$ ,

$$\mathcal{C}^{\mathcal{M},h} = \{x \in \mathcal{C}^0([a, b]) \mid \exists t_0 \in [a, b] : \forall t \in [a, b] :$$

$$x \text{ not smooth at } t \Rightarrow t - t_0 \in h\mathbb{N}\}.$$



Classical variational problem



Meshed variational problem



# The meshed variational problem

Criticality conditions of a meshed variational problem:

Euler-Lagrange equations:  $\frac{\delta \mathcal{L}}{\delta x} = 0$ ,



Natural interior conditions:  $\forall j \geq 2 : \frac{\delta \mathcal{L}}{\delta x^{(j)}} = 0$ ,

or equivalently:  $\forall j \geq 2 : \frac{\partial \mathcal{L}}{\partial x^{(j)}} = 0$ ,

where

$$\frac{\delta \mathcal{L}}{\delta x^{(j)}} = \sum_{k=0}^{\infty} (-1)^k \frac{d^k}{dt^k} \frac{\partial \mathcal{L}}{\partial x^{(j+k)}}.$$

If  $\mathcal{L}$  is a non-convergent power series, these equations are formal.



## The meshed variational problem

We have that  $S_{\text{disc}}(x(0), x(h), \dots) = \int \mathcal{L}_{\text{mesh}}[x] dt$ .

Hence the discrete action is critical if and only if  $\frac{\delta \mathcal{L}_{\text{mesh}}}{\delta x} = 0$ .

Variations that are supported on a single mesh interval do not change the discrete action, so they cannot change  $\int \mathcal{L}_{\text{mesh}}[x] dt$ .

Since these are the variations that produce natural interior conditions, it follows that they are automatically satisfied:

$$\frac{\delta \mathcal{L}_{\text{mesh}}}{\delta x} = 0 \quad \Rightarrow \quad \forall j \geq 2 : \frac{\partial \mathcal{L}}{\partial x^{(j)}} = 0$$

## The meshed variational problem

We have that  $S_{\text{disc}}(x(0), x(h), \dots) = \int \mathcal{L}_{\text{mesh}}[x] dt$ .

Hence the discrete action is critical if and only if  $\frac{\delta \mathcal{L}_{\text{mesh}}}{\delta x} = 0$ .

Variations that are supported on a single mesh interval do not change the discrete action, so they cannot change  $\int \mathcal{L}_{\text{mesh}}[x] dt$ .

Since these are the variations that produce natural interior conditions, it follows that they are automatically satisfied:

$$\frac{\delta \mathcal{L}_{\text{mesh}}}{\delta x} = 0 \quad \Rightarrow \quad \forall j \geq 2 : \frac{\partial \mathcal{L}}{\partial x^{(j)}} = 0$$

The modified equation can be calculated as

$$\frac{\partial \mathcal{L}_{\text{mesh}}}{\partial x} - \frac{d}{dt} \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial \dot{x}} = 0$$

even though the Lagrangian  $\mathcal{L}_{\text{mesh}}$  depends on higher derivatives.

# Table Of Contents

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]

- [1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.
- [2] V. [Modified equations for variational integrators](#). Numer. Math. 137:1001, 2017.
- [3] V. [Numerical precession in variational discretizations of the Kepler problem](#). arXiv:1602.01049.
- [4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). arXiv:1709.09567.

## Definition

The *modified Lagrangian* is the formal power series

$$\mathcal{L}_{\text{mod}}(x, \dot{x}) = \mathcal{L}_{\text{mesh}}[x] \Big|_{\ddot{x}=f_h(x, \dot{x}), \ x^{(3)}=\frac{d}{dt}f_h(x, \dot{x}), \ \dots},$$

where  $\ddot{x} = f_h(x, \dot{x})$  is the modified equation.

The  $k$ -th truncation of the modified Lagrangian is

$$\mathcal{L}_{\text{mod},k} = \mathcal{T}_k(\mathcal{L}_{\text{mod}}(x, \dot{x})) = \mathcal{T}_k\left(\mathcal{L}_{\text{mesh}}[x] \Big|_{x^{(j)}=F_{k-2}^j(x, \dot{x})}\right),$$

where  $\mathcal{T}_k$  denotes truncation after the  $h^k$ -term and

$$\begin{aligned}\ddot{x} &= F_k^2(x, \dot{x}; h) + \mathcal{O}(h^{k+1}) = F_k(x, \dot{x}; h) + \mathcal{O}(h^{k+1}), \\ x^{(3)} &= F_k^3(x, \dot{x}; h) + \mathcal{O}(h^{k+1}), \quad x^{(4)} = F_k^4(x, \dot{x}; h) + \mathcal{O}(h^{k+1}), \quad \dots\end{aligned}$$

are the  $k$ -th truncation of the modified equation and its derivatives.

## Lemma

The meshed modified Lagrangian  $\mathcal{L}_{\text{mesh}}[x]$  and the modified Lagrangian  $\mathcal{L}_{\text{mod}}(x, \dot{x})$  have the same critical curves.

Proof.

$$\begin{aligned} \frac{\partial \mathcal{L}_{\text{mod},k}}{\partial x} &= \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial x} + \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial \ddot{x}} \frac{\partial F_k^2}{\partial x} + \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial x^{(3)}} \frac{\partial F_k}{\partial x} + \dots \Big|_{x^{(j)}=F_{k-1}^j(x, \dot{x})} \\ &= \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial x} + \mathcal{O}(h^{k+1}), \end{aligned}$$

Also,

$$\begin{aligned} \frac{\partial \mathcal{L}_{\text{mod},k}}{\partial \dot{x}} &= \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial \dot{x}} + \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial \ddot{x}} \frac{\partial F_k^2}{\partial \dot{x}} + \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial x^{(3)}} \frac{\partial F_k^3}{\partial \dot{x}} + \dots \Big|_{x^{(j)}=F_{k-1}^j(x, \dot{x})} \\ &= \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial \dot{x}} + \mathcal{O}(h^{k+1}), \\ \Rightarrow \frac{\partial \mathcal{L}_{\text{mod},k}}{\partial x} - \frac{d}{dt} \frac{\partial \mathcal{L}_{\text{mod},k}}{\partial \dot{x}} &= \sum_{j=0}^{\infty} (-1)^j \frac{d^j}{dt^j} \frac{\partial \mathcal{L}_{\text{mesh}}}{\partial x^{(j)}} + \mathcal{O}(h^{k+1}). \end{aligned}$$



# Main result

## Theorem

*For a discrete Lagrangian  $L_{\text{disc}}$  that is a consistent discretization of some  $\mathcal{L}$ , the  $k$ -th truncation of the Euler-Lagrange equation of  $\mathcal{L}_{\text{mod},k}(x, \dot{x})$  is the  $k$ -th truncation of the modified equation.*

**Proof.** Let  $x$  be a solution of the Euler-Lagrange equation for  $\mathcal{L}_{\text{mod}}(x, \dot{x})$ . Consider the discrete curve  $x_j = x(jh)$ .

- ▶  $x$  is critical for the action  $\int \mathcal{L}_{\text{mod}}(x, \dot{x}) dt$ .
- ▶ By the Lemma,  $x$  is critical for the action  $\int \mathcal{L}_{\text{mesh}}[x] dt$ .
- ▶ By construction, the actions  $S_{\text{disc}} = \sum_j L_{\text{disc}}(y(jh), y((j+1)h))$  and  $S = \int_a^b \mathcal{L}_{\text{mod}}[y(t)] dt$  are equal for any smooth curve  $y$ .
- ▶ Therefore the discrete curve  $(x(jh))_j$  is critical for the discrete action  $S_{\text{disc}}$ . Hence

$$D_2 L_{\text{disc}}(x(t-h), x(t)) + D_1 L_{\text{disc}}(x(t), x(t+h)) = 0. \quad \blacksquare$$

# Table Of Contents

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]

- [1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.
- [2] V. [Modified equations for variational integrators](#). Numer. Math. 137:1001, 2017.
- [3] V. [Numerical precession in variational discretizations of the Kepler problem](#). arXiv:1602.01049.
- [4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). arXiv:1709.09567.



## Example: Störmer-Verlet discretization

$$\mathcal{L}(x, \dot{x}) = \frac{1}{2} |\dot{x}|^2 - U(x),$$

$$L_{\text{disc}}(x_j, x_{j+1}) = \frac{1}{2} \left| \frac{x_{j+1} - x_j}{h} \right|^2 - \frac{1}{2} U(x_j) - \frac{1}{2} U(x_{j+1}).$$

Its Euler-Lagrange equation is

$$\frac{x_{j+1} - 2x_j + x_{j-1}}{h^2} = -U'(x_j).$$

We have

$$\begin{aligned} \mathcal{L}_{\text{disc}}[x] &= \left\langle \dot{x} + \frac{h^2}{24} x^{(3)} + \dots, \dot{x} + \frac{h^2}{24} x^{(3)} + \dots \right\rangle \\ &\quad - \frac{1}{2} U \left( x - \frac{h}{2} \dot{x} + \frac{1}{2} \left( \frac{h}{2} \right)^2 \ddot{x} - \dots \right) - \frac{1}{2} U \left( x + \frac{h}{2} \dot{x} + \frac{1}{2} \left( \frac{h}{2} \right)^2 \ddot{x} + \dots \right) \\ &= \frac{1}{2} |\dot{x}|^2 - U + \frac{h^2}{24} \left( \langle \dot{x}, x^{(3)} \rangle - 3U' \ddot{x} - 3U''(\dot{x}, \dot{x}) \right) + \mathcal{O}(h^4) \end{aligned}$$

## Example: Störmer-Verlet discretization

$$\mathcal{L}_{\text{disc}}[x] = \frac{1}{2}|\dot{x}|^2 - U + \frac{h^2}{24} \left( \langle \dot{x}, x^{(3)} \rangle - 3U'\ddot{x} - 3U''(\dot{x}, \dot{x}) \right) + \mathcal{O}(h^4),$$

From this we calculate the meshed modified Lagrangian,

$$\begin{aligned}\mathcal{L}_{\text{mesh}}[x] &= \mathcal{L}_{\text{disc}}[x] - \frac{h^2}{24} \frac{d^2}{dt^2} \mathcal{L}_{\text{disc}}[x] + \mathcal{O}(h^4) \\ &= \frac{1}{2}|\dot{x}|^2 - U + \frac{h^2}{24} \left( \langle \dot{x}, x^{(3)} \rangle - 3U'\ddot{x} - 3U''(\dot{x}, \dot{x}) \right) \\ &\quad - \frac{h^2}{24} \left( \langle \ddot{x}, \ddot{x} \rangle + \langle \dot{x}, x^{(3)} \rangle - U'\ddot{x} - U''(\dot{x}, \dot{x}) \right) + \mathcal{O}(h^4) \\ &= \frac{1}{2}|\dot{x}|^2 - U + \frac{h^2}{24} \left( -\langle \ddot{x}, \ddot{x} \rangle - 2U'\ddot{x} - 2U''(\dot{x}, \dot{x}) \right) + \mathcal{O}(h^4).\end{aligned}$$

Eliminate second derivatives using  $\ddot{x} = -U' + \mathcal{O}(h^2)$ ,

$$\mathcal{L}_{\text{mod},3}(x, \dot{x}) = \frac{1}{2}|\dot{x}|^2 - U + \frac{h^2}{24} (|U'|^2 - 2U''(\dot{x}, \dot{x})).$$

## Example: Störmer-Verlet discretization

The modified Lagrangian is

$$\mathcal{L}_{\text{mod},3}(x, \dot{x}) = \frac{1}{2} \langle \dot{x}, \dot{x} \rangle - U + \frac{h^2}{24} (|U'|^2 - 2U''(\dot{x}, \dot{x})) .$$

Observe that this Lagrangian is not separable for general  $U$ .

The corresponding Euler-Lagrange equation is

$$-\ddot{x} - U' + \frac{h^2}{24} (2U''U' - 2U'''(\dot{x}, \dot{x}) + 4U'''(\dot{x}, \dot{x}) + 4U''\ddot{x}) = 0.$$

Solving this for  $\ddot{x}$  we find the modified equation

$$\ddot{x} = -U' + \frac{h^2}{12} (U'''(\dot{x}, \dot{x}) - U''U') + \mathcal{O}(h^4).$$

# The Kepler problem


Potential:  $U(x) = -\frac{1}{|x|}$ .

Lagrangian:  $\mathcal{L} = \frac{1}{2} \langle \dot{x}, \dot{x} \rangle + \frac{1}{|x|}$ .

Equation of motion  $\ddot{x} = -\frac{x}{|x|^3}$ .

Störmer-Verlet discretization:

$$\frac{x_{j+1} - 2x_j + x_{j-1}}{h^2} = -U'(x_j).$$



# The Kepler problem

Potential:  $U(x) = -\frac{1}{|x|}$ .

Lagrangian:  $\mathcal{L} = \frac{1}{2} \langle \dot{x}, \dot{x} \rangle + \frac{1}{|x|}$ .


Equation of motion  $\ddot{x} = -\frac{x}{|x|^3}$ .

Störmer-Verlet discretization:

$$\frac{x_{j+1} - 2x_j + x_{j-1}}{h^2} = -U'(x_j).$$

Midpoint discretization:

$$\frac{x_{j+1} - 2x_j + x_{j-1}}{h^2} = -\frac{1}{2}U'\left(\frac{x_{j-1} + x_j}{2}\right) - \frac{1}{2}U'\left(\frac{x_j + x_{j+1}}{2}\right).$$



# Störmer-Verlet discretization of the Kepler problem

The modified Lagrangian of the Störmer-Verlet discretization is

$$\mathcal{L}_{\text{mod},3}(x, \dot{x}) = \frac{1}{2}|\dot{x}|^2 - U + \frac{h^2}{24} (U' U' - 2U''(\dot{x}, \dot{x})) .$$

For the Kepler problem we have  $U(x) = -\frac{1}{|x|}$ , hence

$$\mathcal{L}_{\text{mod},3}(x, \dot{x}) = \frac{1}{2}|\dot{x}|^2 + \frac{1}{|x|} + \frac{h^2}{24} \left( \frac{1}{|x|^4} - 2\frac{\langle \dot{x}, \dot{x} \rangle}{|x|^3} + 6\frac{\langle x, \dot{x} \rangle^2}{|x|^5} \right) .$$

# Perturbation theory

The direction and shape of an elliptic orbit is determined by the Laplace-Runge-Lenz vector, which is the Noether integral for a generalized variational symmetry.

Introducing perturbations into Noether's theorem we find

## Lemma

*The precession rate (in radians per period) for the perturbed Lagrangian*

$$\mathcal{L} = \frac{1}{2} \langle \dot{\mathbf{x}}, \dot{\mathbf{x}} \rangle + \frac{1}{|\mathbf{x}|} + \Delta U(\mathbf{x}, \dot{\mathbf{x}}),$$

*is given in first order approximation by*

$$2\pi a^2 \frac{\partial \langle \Delta U(\mathbf{x}, \dot{\mathbf{x}}) \rangle}{\partial b},$$



*where  $a$  and  $b$  are the semimajor and semiminor axes of the orbit respectively, and  $\langle \cdot \rangle$  denotes the time-average along the unperturbed orbit.*

# Störmer-Verlet discretization of the Kepler problem

## Proposition

*The numerical precession rate of the Störmer-Verlet method is*

$$\frac{\pi}{24} \left( 15 \frac{a^3}{b^6} - 3 \frac{a}{b^4} \right) h^2 + \mathcal{O}(h^4)$$



Predicted:  
0.0673 rad per  
revolution.

Measured:  
0.0659 rad per  
revolution.





# Midpoint discretization of the Kepler problem

## Proposition

*The numerical precession rate of the midpoint rule is*

$$-\frac{\pi}{12} \left( 15 \frac{a^3}{b^6} - 3 \frac{a}{b^4} \right) h^2 + \mathcal{O}(h^4)$$



Predicted:  
−0.134 rad per  
revolution.

Measured:  
−0.152 rad per  
revolution.

## New methods

$$\text{Störmer-Verlet: } \frac{\pi}{24} \left( 15 \frac{a^3}{b^6} - 3 \frac{a}{b^4} \right) h^2 + \mathcal{O}(h^4)$$

$$\text{Midpoint rule: } -\frac{\pi}{12} \left( 15 \frac{a^3}{b^6} - 3 \frac{a}{b^4} \right) h^2 + \mathcal{O}(h^4)$$

This allows us to construct new integrators with precession of order  $h^4$ .

(Related idea: Chartier, Hairer, Vilmart. [Numerical integrators based on modified differential equations](#), 2007.)

### Mixed Lagrangian (ML)

$$L(x_j, x_{j+1}) = \frac{2}{3} L_{SV}(x_j, x_{j+1}) + \frac{1}{3} L_{MP}(x_j, x_{j+1})$$

Produces an implicit method, given by

$$x_{j+1} - 2x_j + x_{j-1} = -\frac{2h^2}{3} U'(x_j) - \frac{h^2}{6} U' \left( \frac{x_{j-1} + x_j}{2} \right) - \frac{h^2}{6} U' \left( \frac{x_j + x_{j+1}}{2} \right)$$

# New methods

## Lagrangian composition (LC)

Consider the discrete Lagrangians

$$L_j(x_k, x_{k+1}) = \begin{cases} L_{MP}(x_k, x_{k+1}) & \text{if } 3|j, \\ L_{SV}(x_k, x_{k+1}) & \text{otherwise.} \end{cases}$$

Three different Euler-Lagrange equations which are applied for different values of  $j \bmod 3$ :

$$\begin{cases} x_{j+1} - 2x_j + x_{j-1} = -\frac{h^2}{2} U' \left( \frac{x_{j-1} + x_j}{2} \right) - \frac{h^2}{2} U'(x_j) & \text{if } j \equiv 0 \bmod 3, \\ x_{j+1} - 2x_j + x_{j-1} = -h^2 U'(x_j) & \text{if } j \equiv 1 \bmod 3, \\ x_{j+1} - 2x_j + x_{j-1} = -\frac{h^2}{2} U' \left( \frac{x_j + x_{j+1}}{2} \right) - \frac{h^2}{2} U'(x_j) & \text{if } j \equiv 2 \bmod 3. \end{cases}$$

Equivalent to composing the corresponding symplectic maps.


# New methods

## Composition of difference equations (DEC)


$$\begin{cases} x_{j+1} - 2x_j + x_{j-1} = -\frac{h^2}{2} U' \left( \frac{x_{j-1} + x_j}{2} \right) - \frac{h^2}{2} U' \left( \frac{x_j + x_{j+1}}{2} \right) & \text{if } j \equiv 2 \pmod{3}, \\ x_{j+1} - 2x_j + x_{j-1} = -h^2 U'(x_j) & \text{otherwise.} \end{cases}$$

Is this still a variational integrator?

For any of the new methods:



# Precession rates



MP,SV: old methods

LC, ML, DEC: new methods

FR: Forest, Ruth. [Fourth-order symplectic integration](#), 1989.

C: Chin. [Symplectic integrators from composite operator factorizations](#), 1997.

# Table Of Contents

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]

[1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.

[2] V. [Modified equations for variational integrators](#). Numer. Math. 137:1001, 2017.

[3] V. [Numerical precession in variational discretizations of the Kepler problem](#).  
arXiv:1602.01049.

[4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). arXiv:1709.09567.

# Lagrangians linear in velocities

$\mathcal{L} : T\mathbb{R}^N \cong \mathbb{R}^{2N} \rightarrow \mathbb{R}$  of the form

$$\mathcal{L}(q, \dot{q}) = \langle \alpha(q), \dot{q} \rangle - H(q),$$

where  $\alpha : \mathbb{R}^N \rightarrow \mathbb{R}^N$ ,  $H : \mathbb{R}^N \rightarrow \mathbb{R}$ , and the brackets  $\langle \cdot, \cdot \rangle$  denote the standard scalar product.

Let

$$A(q) = \alpha'(q) = \left( \frac{\partial \alpha_i(q)}{\partial q_j} \right)_{i,j=1,\dots,N} \quad \text{and} \quad A_{\text{skew}}(q) = A(q)^T - A(q)$$

We assume that  $A_{\text{skew}}(q)$  is invertible, then the Euler-Lagrange equation for  $\mathcal{L}$  is given by

$$\dot{q} = A_{\text{skew}}(q)^{-1} H'(q)^T$$

# Examples of Lagrangians linear in velocities

- Dynamics of point vortices in the (complex) plane

$$\mathcal{L}(z, \bar{z}, \dot{z}, \dot{\bar{z}}) = \sum_{j=1}^N \Gamma_j \operatorname{Im}(\bar{z}_j \dot{z}_j) - \frac{1}{\pi} \sum_{j=1}^N \sum_{k=1}^{j-1} \Gamma_j \Gamma_k \log |z_j - z_k|,$$

$$\hookrightarrow \quad \dot{z}_j = \frac{i}{2\pi} \sum_{k \neq j} \frac{\Gamma_k}{\bar{z}_j - \bar{z}_k} \quad \text{for } j = 1, \dots, N.$$

- Variational formulation in phase space

$$\mathcal{L}(p, q, \dot{p}, \dot{q}) = \langle p, \dot{q} \rangle - H(p, q).$$

$$\hookrightarrow \quad \dot{q} = \left( \frac{\partial H}{\partial p} \right)^T \quad \text{and} \quad \dot{p} = - \left( \frac{\partial H}{\partial q} \right)^T.$$

- Guiding centre motion (plasma physics)
- Many PDEs, e.g. nonlinear Schrödinger equation.

(But modified equations are not so useful for PDEs)



## Possible discretization of $\mathcal{L}(q, \dot{q}) = \langle \alpha(q), \dot{q} \rangle - H(q)$

$$L_{\text{disc}}(q_j, q_{j+1}, h) = \left\langle \frac{1}{2}\alpha(q_j) + \frac{1}{2}\alpha(q_{j+1}), \frac{q_{j+1} - q_j}{h} \right\rangle - \frac{1}{2}H(q_j) - \frac{1}{2}H(q_{j+1})$$

$$\hookrightarrow \left( \frac{q_{j+1} - q_{j-1}}{2h} \right)^T \alpha'(q_j) - \frac{\alpha(q_{j+1})^T - \alpha(q_{j-1})^T}{2h} - H'(q_j) = 0.$$

In case  $\alpha$  is linear the Euler-Lagrange equation simplifies to


$$\frac{q_{j+1} - q_{j-1}}{2h} = A_{\text{skew}}^{-1} H'(q_j)^T.$$

The EL equation involves 3 points  $\Rightarrow$  needs 2 points of initial data.

The differential equation is of 1st order  $\Rightarrow$  needs only 1 point of initial data.

This means we are dealing with a 2-step method and parasitic solutions can occur.

# Parasitic solutions



Every multi-step method has an underlying 1-step method.

If the initial data lie on a solution of this 1-step method, both will agree.

If not, the solution of the multistep method oscillates around the solutions of the 1-step method. These oscillations can be bounded or exponentially growing, depending on the method.

In case the oscillations grow, parasitic oscillations take over after a certain time.

Even with perfect initial data, rounding errors will introduce oscillations.

# Modified equations for 2-step methods

## Principal modified equation

$$\dot{q} = f(q) + hf_1(q) + h^2f_2(q) + \dots + h^kf_k(q)$$

satisfies

$$\begin{aligned} & \frac{a_0q(t) + a_1q(t+h) + a_2q(t+2h)}{h} \\ &= b_0f(q(t)) + b_1f(q(t+h)) + b_2f(q(t+2h)) + \mathcal{O}(h^{k+1}). \end{aligned}$$

## Full system of modified equations

$$\begin{aligned} \dot{x} &= f_0(x, y) + hf_1(x, y) + \dots + h^kf_k(x, y) \\ \dot{y} &= g_0(x, y) + hg_1(x, y) + \dots + h^kg_k(x, y), \end{aligned}$$

such that the discrete curve  $q_j = x(t+jh) + (-1)^j y(t+jh)$  satisfies

$$\frac{a_0q_j + a_1q_{j+1} + a_2q_{j+2}}{h} = b_0f(q_j) + b_1f(q_{j+1}) + b_2f(q_{j+2}) + \mathcal{O}(h^{k+1})$$

# A Lagrangian for the principal modified equation

Exactly the same as in the non-degenerate case:

- 1 Taylor expansion to get  $\mathcal{L}_{\text{disc}}$ ,
- 2 Euler-Maclaurin formula to get  $\mathcal{L}_{\text{mesh}}$ ,
- 3 Replace higher derivatives to get  $\mathcal{L}_{\text{mod}}$ .

Even though we now have a first-order equation, we still **cannot replace first derivatives** in the Lagrangian.

Replacement of derivatives is allowed because of the natural interior conditions,

$$\forall \ell \geq 2 : \quad \frac{\partial \mathcal{L}}{\partial q^{(\ell)}}(t) = 0.$$

## Doubling the dimension

The discrete curve  $(x_j, y_j)_{j \in \mathbb{Z}}$  is critical for

$$\widehat{L}(x_j, y_j, x_{j+1}, y_{j+1}, h) = \frac{1}{2}L(x_j + y_j, x_{j+1} - y_{j+1}, h) + \frac{1}{2}L(x_j - y_j, x_{j+1} + y_{j+1}, h),$$

if and only if the discrete curves  $(q_j^+)_{j \in \mathbb{Z}}$  and  $(q_j^-)_{j \in \mathbb{Z}}$ , defined by

$$q_j^\pm = x_j \pm (-1)^j y_j,$$

are critical for  $L(q_j, q_{j+1}, h)$ .

Lagrangian for the full system of modified equations

= Lagrangian for the principal modified equation of the extended system.

Hence we can calculate a Lagrangian for the full system of modified equations with the tools we already have.

## Example 1

For

$$L_{\text{disc}}(q_j, q_{j+1}, h) = \left\langle \frac{1}{2}Aq_j + \frac{1}{2}Aq_{j+1}, \frac{q_{j+1} - q_j}{h} \right\rangle - \frac{1}{2}H(q_j) - \frac{1}{2}H(q_{j+1})$$

we find

$$\widehat{\mathcal{L}}_{\text{mod},0}(x, y, \dot{x}, \dot{y}, h) = \langle Ax, \dot{x} \rangle + \langle A\dot{y}, y \rangle - \frac{1}{2}H(x+y) - \frac{1}{2}H(x-y).$$

Its Euler-Lagrange equations are

$$\begin{aligned}\dot{x} &= A_{\text{skew}}^{-1} \left( \frac{1}{2}H'(x+y)^T + \frac{1}{2}H'(x-y)^T \right) + \mathcal{O}(h), \\ \dot{y} &= A_{\text{skew}}^{-1} \left( -\frac{1}{2}H'(x+y)^T + \frac{1}{2}H'(x-y)^T \right) + \mathcal{O}(h).\end{aligned}$$

Linearize the second equation around  $y = 0$

$$\dot{y} = -A_{\text{skew}}^{-1}H''(x)y + \mathcal{O}(|y|^2 + h)$$


# Example 1

Magnitude of oscillations satisfies

$$\dot{y} = -A_{\text{skew}}^{-1} H''(x)y + \mathcal{O}(|y|^2 + h)$$

Unless the matrix  $-A_{\text{skew}}^{-1} H''(x)$  is exceptionally friendly, we expect growing parasitic oscillations.

(Note that an eigenvalue analysis does not apply because  $-A_{\text{skew}}^{-1} H''(x)$  is not constant)



## Example 2

For

$$L_{\text{disc}}(q_j, q_{j+1}, h) = \left\langle A \frac{q_j + q_{j+1}}{2}, \frac{q_{j+1} - q_j}{h} \right\rangle - H\left(\frac{q_j + q_{j+1}}{2}\right)$$


we find

$$\hat{\mathcal{L}}_{\text{mod},0}(x, y, \dot{x}, \dot{y}, h) = \langle A x, \dot{x} \rangle + \langle A y, \dot{y} \rangle - H(x).$$

Its Euler-Lagrange equations are

$$\begin{aligned}\dot{x} &= A_{\text{skew}}^{-1} H'(x)^T + \mathcal{O}(h), \\ \dot{y} &= 0 + \mathcal{O}(h).\end{aligned}$$

Even better,  $\dot{y} = 0$  to any order  $\rightarrow$  no growing oscillations.





# Summary

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]

- [1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.
- [2] V. [Modified equations for variational integrators](#). *Numer. Math.* 137:1001, 2017.
- [3] V. [Numerical precession in variational discretizations of the Kepler problem](#).  
[arXiv:1602.01049](#).
- [4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). [arXiv:1709.09567](#).

# Summary

- ▶ Obtaining a high-order modified Lagrangian  $\mathcal{L}_{\text{mesh}}[x]$  is relatively straightforward, but its interpretation is not.
- ▶ From  $\mathcal{L}_{\text{mesh}}[x]$  a first order Lagrangian  $\mathcal{L}_{\text{mod},k}(x, \dot{x})$  can be found using the meshed variational principle.
- ▶ If the Lagrangian is nondegenerate, the modified Lagrangian can also be obtained by Legendre transform from the modified Hamiltonian.
- ▶ Our approach extends to degenerate Lagrangians that are linear in velocities.
- ▶ Can we get improved error estimates from the Lagrangian perspective?
- ▶ What about nonholonomic constraints?
- ▶ What about PDEs?

# ¡Gracias por su atención!

- 1 Variational and symplectic integrators [1]
- 2 Modified Equations [1]
- 3 A meshed modified Lagrangian [2]
- 4 A true modified Lagrangian [2]
- 5 Variational integration of the Kepler problem [3]
- 6 A class of degenerate Lagrangians [4]

[1] Hairer, Lubich, Wanner. [Geometric Numerical Integration](#). Springer, 2006.

[2] V. [Modified equations for variational integrators](#). Numer. Math. 137:1001, 2017.

[3] V. [Numerical precession in variational discretizations of the Kepler problem](#).  
arXiv:1602.01049.

[4] V. [Modified equations for variational integrators applied to Lagrangians linear in velocities](#). arXiv:1709.09567.