



Loughborough  
University

# Semi-discrete pluri-Lagrangian structures and the Toda hierarchy

Mats Vermeeren

DGD Seminar  
Berlin, 28 June 2022

# Contents

## ● Introduction

### ① Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries

### ② Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy

### ③ Summary

# Liouville-Arnold integrability

A Hamiltonian system with Hamilton function  $H : T^*Q \cong \mathbb{R}^{2N} \rightarrow \mathbb{R}$  is **Liouville integrable** if there exist  $N$  functionally independent Hamilton functions  $H = H_1, H_2, \dots, H_N$  such that  $\{H_i, H_j\} = 0$ .

- ▶ Each  $H_i$  defines its own flow:  $N$  dynamical systems
- ▶ Each  $H_i$  is a conserved quantity for all flows.
- ▶ The dynamics is confined to a leaf of the foliation  $\{H_i = \text{const}\}$ .
- ▶ If this foliation is compact, its leaves are tori.
- ▶ Dynamics on these tori are linear in action-angle variables.
- ▶ **The flows commute:**

$$\phi_{H_i}^t \circ \phi_{H_j}^s = \phi_{H_j}^s \circ \phi_{H_i}^t.$$


(Infinitesimally:  $[X_{H_i}, X_{H_j}] = 0$ .)

## Two commuting flows

Let  $z = (q, p)$ . Consider two Hamiltonian ODEs

$$\begin{aligned}\frac{df(z)}{dt_1} &= \{f(z), H_1(z)\} \\ \frac{df(z)}{dt_2} &= \{f(z), H_2(z)\}\end{aligned}\quad \text{with } \{H_1, H_2\} = 0$$

The flows commute, meaning that evolution can be parameterised by the  $(t_1, t_2)$  plane, called **multi-time**.



Additional commuting equations can be accommodated by increasing the dimension of multi-time:  $\mathbb{R}^N$  instead of  $\mathbb{R}^2$ .

# Lagrangian mechanics

Lagrange function  $L : TQ \cong \mathbb{R}^{2N} \rightarrow \mathbb{R} : (q, q_t) \mapsto L(q, q_t)$

Dynamics follows curves which are minimizers (critical points) of the action

$$\int_a^b L(q, q_t) dt \quad \text{with fixed boundary values } q(a) \text{ and } q(b).$$

Minimizers satisfy the Euler-Lagrange (EL) equation  $\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial q_t} = 0$

**Proof.** Consider an arbitrary variation  $\delta q$ :

$$\delta \int_a^b L dt = \int_a^b \left( \frac{\partial L}{\partial q} \delta q + \frac{\partial L}{\partial q_t} \delta q_t \right) dt$$

Integration by parts yields

$$\delta \int_a^b L dt = \int_a^b \left( \frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial q_t} \right) \delta q dt + \left[ \frac{\partial L}{\partial q_t} \delta q \right]_a^b$$

EL follows because  $\delta q(a) = \delta q(b) = 0$  and  $\delta q$  is arbitrary inside  $(a, b)$ . ■

# Contents

## ● Introduction

### 1 Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries

### 2 Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy

### 3 Summary

Includes work by Nijhoff, Suris, Petrera, Sleight, ...

# Contents

## ● Introduction

### 1 Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries

### 2 Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy

### 3 Summary

# Lagrangian formulation of Liouville integrable system

On the Hamiltonian side, commutativity is implied by  $\{H_i, H_j\} = 0$ .

What about the Lagrangian side?

Suppose we have Lagrange functions  $L_i$  associated to  $H_i$ .

## Variational (“Pluri-Lagrangian”) principle for ODEs


Combine the  $L_i$  into a 1-form

$$\mathcal{L}[q] = \sum_{i=1}^N L_i[q] dt_i.$$

Look for dynamical variables  $q(t_1, \dots, t_N)$  such that the action

$$\mathcal{I}_S = \int_S \mathcal{L}[q]$$

is critical w.r.t. variations of  $q$ , simultaneously over every curve  $S$  in multi-time  $\mathbb{R}^N$



# Multi-time Euler-Lagrange equations

Assume that

$$L_1[q] = L_1(q, q_{t_1}),$$

$$L_i[q] = L_i(q, q_{t_1}, q_{t_i}), \quad i \neq 1$$

The multi-time Euler-Lagrange equations for  $\mathcal{L} = \sum_i L_i[q] dt_i$  are

Usual Euler-Lagrange equations: 
$$\frac{\partial L_i}{\partial q} - \frac{d}{dt_i} \frac{\partial L_i}{\partial q_{t_i}} = 0$$

Usual EL wrt to alien derivatives: 
$$\frac{\partial L_i}{\partial q_{t_1}} - \frac{d}{dt_i} \frac{\partial L_i}{\partial q_{t_1 t_i}} = 0, \quad i \neq 1$$

Additional conditions: 
$$\frac{\partial L_i}{\partial q_{t_i}} = \frac{\partial L_j}{\partial q_{t_j}}$$

## Example: Kepler Problem

The classical Lagrangian

$$L_1[q] = \frac{1}{2}|q_{t_1}|^2 + \frac{1}{|q|}$$

can be combined with

$$L_2[q] = q_{t_1} \cdot q_{t_2} + (q_{t_1} \times q) \cdot e \quad (e \text{ fixed unit vector})$$

into a Lagrangian 1-form  $\mathcal{L} = L_1 dt_1 + L_2 dt_2$ .

Multi-time Euler-Lagrange equations:

$$\frac{\partial L_1}{\partial q} - \frac{d}{dt_1} \frac{\partial L_1}{\partial q_{t_1}} = 0 \quad \Rightarrow \quad q_{t_1 t_1} = -\frac{q}{|q|^3} \quad (\text{Keplerian motion})$$

$$\frac{\partial L_2}{\partial q} - \frac{d}{dt_2} \frac{\partial L_2}{\partial q_{t_2}} = 0 \quad \Rightarrow \quad q_{t_1 t_2} = e \times q_{t_1}$$

$$\frac{\partial L_2}{\partial q_{t_1}} = 0 \quad \Rightarrow \quad q_{t_2} = e \times q \quad (\text{Rotation})$$

$$\frac{\partial L_1}{\partial q_{t_1}} = \frac{\partial L_2}{\partial q_{t_2}} \quad \Rightarrow \quad q_{t_1} = q_{t_1}$$

# Derivation of the multi-time Euler-Lagrange equations


Consider a Lagrangian one-form  $\mathcal{L} = \sum_i L_i[q] dt_i$ , with

$$L_1[q] = L_1(q, q_{t_1}),$$


$$L_i[q] = L_i(q, q_{t_1}, q_{t_i}), \quad i \neq 1$$

## Lemma

If the action  $\int_S \mathcal{L}$  is critical on all **stepped curves**  $S$  in  $\mathbb{R}^N$ , then it is critical on all smooth curves.



Variations are local, so it is sufficient to look at an **L-shaped curve**  $S = S_i \cup S_j$ .



# Derivation of the multi-time Euler-Lagrange equations

On one of the straight pieces,  $S_i$  ( $i \neq 1$ ), we get


$$\delta \int_{S_i} L_i dt_i = \int_{S_i} \left( \frac{\partial L_i}{\partial q} \delta q + \frac{\partial L_i}{\partial q_{t_1}} \delta q_{t_1} + \frac{\partial L_i}{\partial q_{t_i}} \delta q_{t_i} \right) dt_i$$

Integration by parts (wrt  $t_i$  only) yields

$$\delta \int_{S_i} L_i dt_i = \int_{S_i} \left( \left( \frac{\partial L_i}{\partial q} - \frac{d}{dt_i} \frac{\partial L_i}{\partial q_{t_i}} \right) \delta q + \frac{\partial L_i}{\partial q_{t_1}} \delta q_{t_1} \right) dt_i + \frac{\partial L_i}{\partial q_{t_i}} \delta q \Big|_C$$

Since  $p$  is an interior point of the curve, we cannot set  $\delta q(C) = 0$ !

Arbitrary  $\delta q$  and  $\delta q_{t_1}$ , so we find:



## Multi-time Euler-Lagrange equations

$$\frac{\partial L_i}{\partial q} - \frac{d}{dt_i} \frac{\partial L_i}{\partial q_{t_i}} = 0, \quad \frac{\partial L_i}{\partial q_{t_1}} - \frac{d}{dt_i} \frac{\partial L_i}{\partial q_{t_1 t_i}} = 0, \quad \frac{\partial L_i}{\partial q_{t_i}} = \frac{\partial L_j}{\partial q_{t_j}}$$

## Higher order Lagrangians $L_i[q] = L_i(q, q_{t_i}, q_{t_i t_i}, \dots)$

For a string  $I = t_{i_1} \dots t_{i_k}$  of time variables, denote the corresponding derivative by  $q_I$ .

If  $I$  is empty then  $q_I = q$ .

Denote by  $\frac{\delta_i}{\delta q_I}$  the variational derivative in the direction of  $t_i$  wrt  $q_I$ :

$$\begin{aligned}\frac{\delta_i L_i}{\delta q_I} &= \sum_{\alpha=0}^{\infty} (-1)^\alpha \frac{d^\alpha}{dt_i^\alpha} \frac{\partial L_i}{\partial q_{I t_i^\alpha}} \\ &= \frac{\partial L_i}{\partial q_I} - \frac{d}{dt_i} \frac{\partial L_i}{\partial q_{I t_i}} + \frac{d^2}{dt_i^2} \frac{\partial L_i}{\partial q_{I t_i^2}} - \dots\end{aligned}$$

### Multi-time Euler-Lagrange equations

$$\text{Usual Euler-Lagrange equations: } \frac{\delta_i L_i}{\delta q_I} = 0 \quad \forall I \not\ni t_i,$$

$$\text{Additional conditions: } \frac{\delta_i L_i}{\delta q_{I t_i}} = \frac{\delta_j L_j}{\delta q_{I t_j}} \quad \forall I,$$

# Contents

## ● Introduction

### 1 Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries

### 2 Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy

### 3 Summary



# Variational principle for PDEs ( $d = 2$ )

## Pluri-Lagrangian principle

Given a 2-form

$$\mathcal{L}[q] = \sum_{i,j} L_{ij}[q] dt_i \wedge dt_j,$$

find a field  $q : \mathbb{R}^N \rightarrow \mathbb{R}$ , such that  $\int_S \mathcal{L}[q]$  is **critical on all smooth surfaces**  $S$  in multi-time  $\mathbb{R}^N$ , w.r.t. **variations of  $q$** .




# Multi-time EL equations

for  $\mathcal{L}[q] = \sum_{i,j} L_{ij}[q] dt_i \wedge dt_j$

$$\frac{\delta_{ij} L_{ij}}{\delta q_l} = 0 \quad \forall l \not\supseteq t_i, t_j,$$

$$\frac{\delta_{ij} L_{ij}}{\delta q_{lt_j}} = \frac{\delta_{ik} L_{ik}}{\delta q_{lt_k}} \quad \forall l \not\supseteq t_i,$$

$$\frac{\delta_{ij} L_{ij}}{\delta q_{lt_i t_j}} + \frac{\delta_{jk} L_{jk}}{\delta q_{lt_j t_k}} + \frac{\delta_{ki} L_{ki}}{\delta q_{lt_k t_i}} = 0 \quad \forall l.$$



Where

$$\frac{\delta_{ij} L_{ij}}{\delta q_l} = \sum_{\alpha=0}^{\infty} \sum_{\beta=0}^{\infty} (-1)^{\alpha+\beta} \frac{d^{\alpha}}{dt_i^{\alpha}} \frac{d^{\beta}}{dt_j^{\beta}} \frac{\partial L_{ij}}{\partial q_{lt_i^{\alpha} t_j^{\beta}}}$$

## Example: Potential KdV hierarchy

$$u_{t_2} = Q_2 = u_{xxx} + 3u_x^2,$$

$$u_{t_3} = Q_3 = u_{xxxxx} + 10u_x u_{xxx} + 5u_{xx}^2 + 10u_x^3,$$

where we identify  $t_1 = x$ .

The differentiated equations  $u_{xt_i} = \frac{d}{dx} Q_i$  are Lagrangian with

$$L_{12} = \frac{1}{2} u_x u_{t_2} - \frac{1}{2} u_x u_{xxx} - u_x^3,$$

$$L_{13} = \frac{1}{2} u_x u_{t_3} - \frac{1}{2} u_{xxx}^2 + 5u_x u_{xx}^2 - \frac{5}{2} u_x^4.$$

A suitable coefficient  $L_{23}$  of

$$\mathcal{L} = L_{12} dt_1 \wedge dt_2 + L_{13} dt_1 \wedge dt_3 + L_{23} dt_2 \wedge dt_3$$

can be found (nontrivial task!) in the form

$$L_{23} = \frac{1}{2}(u_{t_3} Q_2 - u_{t_2} Q_3) + p_{23}.$$

## Example: Potential KdV hierarchy

- ▶ The equations  $\frac{\delta_{12}L_{12}}{\delta u} = 0$  and  $\frac{\delta_{13}L_{13}}{\delta u} = 0$  yield

$$u_{xt_2} = \frac{d}{dx} Q_2 \quad \text{and} \quad u_{xt_3} = \frac{d}{dx} Q_3.$$

- ▶ The equations  $\frac{\delta_{12}L_{12}}{\delta u_x} = \frac{\delta_{32}L_{32}}{\delta u_{t_3}}$  and  $\frac{\delta_{13}L_{13}}{\delta u_x} = \frac{\delta_{23}L_{23}}{\delta u_{t_2}}$  yield

$$u_{t_2} = Q_2 \quad \text{and} \quad u_{t_3} = Q_3,$$

the evolutionary equations!

- ▶ All other multi-time EL equations are corollaries of these.

# Contents

## ● Introduction

### 1 Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries

### 2 Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy

### 3 Summary

## Exterior derivative of $\mathcal{L}$

Revisit the **Kepler problem**:  $\mathcal{L} = L_1 dt_1 + L_2 dt_2$  with

$$L_1[q] = \frac{1}{2}|q_{t_1}|^2 + \frac{1}{|q|}$$

$$L_2[q] = q_{t_1} \cdot q_{t_2} + (q_{t_1} \times q) \cdot e \quad (e \text{ fixed unit vector})$$

Multi-time Euler-Lagrange equations:

$$q_{t_1 t_1} = -\frac{q}{|q|^3}$$

$$q_{t_2} = e \times q$$

### Coefficient of $d\mathcal{L}$

$$\frac{dL_2}{dt_1} - \frac{dL_1}{dt_2} = \left( q_{t_1 t_1} + \frac{q}{|q|^3} \right) (q_{t_2} - e \times q)$$


Observation (also for PDEs):  $d\mathcal{L}$  often has a “double zero” on solutions.

$d\mathcal{L} = 0$  sets a **Lagrangian multiform** apart from a pluri-Lagrangian system.

# Interpretation of closedness condition

If  $d\mathcal{L} = 0$ , then the action is **invariant wrt variations in geometry**

Deforming the curve (surface) of integration leaves action invariant.



$$\int_{S_1} \mathcal{L} - \int_{S_2} \mathcal{L} = \int_D d\mathcal{L} = 0$$

Recall: before we only took variations of  $q$ .

$d\mathcal{L}$  provides an **alternative derivation of the EL equations**:

WLOG, we can restrict the variational principle to simple closed curves (surfaces) of integration, i.e. boundaries of submanifolds  $S$ . Then

$$\delta \int_{\partial S} \mathcal{L} = \int_S \delta d\mathcal{L},$$

hence the variational principle is equivalent to  $\delta d\mathcal{L} = 0$ .

Multi-time EL equations can be obtained by taking variations of (coefficients of)  $d\mathcal{L}$ .

# Hamiltonian formulation and $d\mathcal{L}$

We can pass between the pluri-Lagrangian and Hamiltonian formalisms for 1-form and 2-forms.

## Lemma ( $d\mathcal{L}$ for 1-forms)

On solutions there holds  $\frac{dL_j}{dt_i} - \frac{dL_i}{dt_j} = \{H_j, H_i\}$ .

It follows that:

## Theorem

The Hamiltonians are in involution if and only if  $d\mathcal{L} = 0$  on solutions.

A similar result holds for 2-forms (and presumably for higher forms)

# Variational Symmetries and $d\mathcal{L}$

$d\mathcal{L} = 0$  expresses that flows are variational symmetries of each other

$$d\left(\sum_i L_i dt_i\right) = 0 \Rightarrow \frac{dL_k}{dt_j} = \frac{dL_j}{dt_k}$$

$\Rightarrow t_j$ -flow changes  $L_k$  by a  $t_k$ -derivative

$$\Rightarrow \partial_j \int_a^b L_k dt_k = \int_a^b \frac{dL_j}{dt_k} dt_k = [L_j]_a^b = \text{const}$$

Adding a constant to the action does not change the dynamics, hence  $\partial_j$  is a **variational symmetry**.

A similar result holds for higher forms.

We can use variational symmetries to construct Lagrangian multiforms.

# Contents

## ● Introduction

### 1 Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries

### 2 Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy

### 3 Summary

Joint work with Duncan Sleight, [arXiv:2204.130633](https://arxiv.org/abs/2204.130633)

# Contents

## ● Introduction

## ① Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries

## ② Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy


## ③ Summary

# Semi-discrete geometry

We consider only 1 discrete direction:  
multi-time is  $\mathbb{Z} \times \mathbb{R}^N$

A **semi-discrete surface** is a collection of surfaces and curves in  $\mathbb{R}^N$ , each assigned a value of  $\mathbb{Z}$

Curves (white) are where the surface (black) jumps to a different value of  $\mathbb{Z}$



$d$ -dimensional semi-discrete submanifold  $S$

$$S = \left( \bigsqcup_{k \in \mathbb{Z}} S_k^{d-1}, \bigsqcup_{k \in \mathbb{Z}} S_k^d \right)$$

$S_k^{d-1}$ : disjoint union of oriented  $(d-1)$ -submanifolds of  $\mathbb{R}^N$

$S_k^d$ : disjoint union of oriented  $d$ -submanifolds of  $\mathbb{R}^N$ .

$d = 2$ : **semi-discrete surface**

$d = 3$ : **semi-discrete volume**

# Semi-discrete geometry

$$\text{Boundary of } S = \left( \bigsqcup_{k \in \mathbb{Z}} S_k^{d-1}, \bigsqcup_{k \in \mathbb{Z}} S_k^d \right)$$

$$\partial S = \left( \bigsqcup_{k \in \mathbb{Z}} -\partial S_k^{d-1}, \bigsqcup_{k \in \mathbb{Z}} \left( \partial S_k^d \sqcup S_k^{d-1} \sqcup -S_{k+1}^{d-1} \right) \right)$$

where the minus sign denotes a change of orientation.

Sign conventions are chosen so that the boundary of a boundary is empty.

Dynamical variables will be (scalar) functions  $q$  of  $\mathbb{Z} \times \mathbb{R}^N$ .

Superscript to emphasise lattice position:

$$q^{[k]} = q(k, t_1, \dots, t_N)$$

$\mathcal{T}$  denotes shift operator:

$$\mathcal{T} q^{[k]} = q^{[k+1]}$$

# Semi-discrete geometry

- semi-discrete  $d$ -form

$$\mathcal{L}[q] = \left( \mathcal{L}^{d-1}[q], \mathcal{L}^d[q] \right)$$

consists of a  $(d-1)$ -form and a  $d$ -form, with coefficients depending on phase space variables.

- The semi-discrete integral over semi-discrete submanifold  $S$

$$\int_S \mathcal{L}[q] = \sum_k \int_{S_k^{d-1}} \mathcal{L}^{d-1}[q^{[k]}] + \sum_k \int_{S_k^d} \mathcal{L}^d[q^{[k]}],$$

- The exterior derivative:

$$\mathbb{d}\mathcal{L} = \left( \Delta(\mathcal{L}^d) - \mathbb{d}\mathcal{L}^{d-1}, \mathbb{d}\mathcal{L}^d \right),$$

where  $\Delta = \text{id} - \mathcal{T}^{-1}$  is the backward difference operator.

- Stokes theorem:

$$\int_S \mathbb{d}\mathcal{L} = \int_{\partial S} \mathcal{L}.$$

# Variational principle in semi-discrete multi-time


Consider a semi-discrete 2-form

$$\left( \sum_j L_{0j} dt_j, \sum_{i,j} L_{ij} dt_i \wedge dt_j \right)$$

Look for dynamical variables  $q(k, t_1, \dots, t_N)$  such that the action

$$\mathcal{I}_S = \int_S \mathcal{L}[q]$$

is critical w.r.t. **variations of  $q$** , simultaneously over **every semi-discrete surface  $S$**  in multi-time  $\mathbb{R}^N$



# Semi-discrete variational derivatives

$$\frac{\delta_0 L}{\delta q_I^{[k]}} := \frac{\partial}{\partial q_I^{[k]}} \sum_{n \in \mathbb{N}} \mathcal{T}^{-n} L,$$

$$\frac{\delta_{0i} L}{\delta q_I^{[k]}} := \frac{\delta_i}{\delta q_I^{[k]}} \sum_{n \in \mathbb{N}} \mathcal{T}^{-n} L.$$

Traditional discrete EL eqn:

$$\frac{\partial}{\partial q_I^{[k]}} \sum_{n \in \mathbb{Z}} \mathcal{T}^{-n} L = 0$$

Traditional semi-discrete EL eqn:

$$\frac{\delta_i}{\delta q_I^{[k]}} \sum_{n \in \mathbb{Z}} \mathcal{T}^{-n} L = 0$$

Same if  $L$  only depends on  $q_J^{[\ell]}$  for  $\ell \geq k$ .

Denote  $\bar{q} = \mathcal{T}q$  and  $\underline{q} = \mathcal{T}^{-1}q$ .

Examples:

$$\frac{\delta_{0i} q_{t_i}^2}{\delta q} = \frac{\delta_i q_{t_i}^2}{\delta q} = -2 \frac{d}{dt_i} q_{t_i} = -2 q_{t_i t_i},$$

$$\frac{\delta_{0i} \bar{q}_{t_i}^2}{\delta q} = -2 \frac{d}{dt_i} q_{t_i} = -2 q_{t_i t_i},$$

$$\frac{\delta_{0i} q \bar{q}}{\delta q} = \frac{\delta_0 q \bar{q}}{\delta q} = \bar{q} + \underline{q},$$

$$\frac{\delta_{0i} q q}{\delta q} = \frac{\delta_0 q q}{\delta q} = \underline{q}.$$

# Semi-discrete multi-time Euler-Lagrange equations

A field is critical if and only if the following **multi-time Euler-Lagrange equations** hold for all  $n \in \mathbb{Z}$ :

$$\frac{\delta_{ij} L_{ij}}{\delta q_i^{[n]}} = 0 \quad \forall I \not\ni t_i, t_j,$$

$$\frac{\delta_{ij} L_{ij}}{\delta q_{It_j}^{[n]}} - \frac{\delta_{ik} L_{ik}}{\delta q_{It_k}^{[n]}} = 0 \quad \forall I \not\ni t_i,$$

$$\frac{\delta_{ij} L_{ij}}{\delta q_{It_i t_j}^{[n]}} + \frac{\delta_{jk} L_{jk}}{\delta q_{It_j t_k}^{[n]}} + \frac{\delta_{ki} L_{ki}}{\delta q_{It_k t_i}^{[n]}} = 0 \quad \forall I,$$

$$\frac{\delta_{ij} L_{ij}}{\delta q_{It_j}^{[n]}} + \frac{\delta_{0i} L_{0i}}{\delta q_i^{[n]}} = 0 \quad \forall I \not\ni t_i,$$

$$\frac{\delta_{ij} L_{ij}}{\delta q_{It_i t_j}^{[n]}} - \frac{\delta_{0j} L_{0j}}{\delta q_{It_j}^{[n]}} + \frac{\delta_{0i} L_{0i}}{\delta q_{It_i}^{[n]}} = 0 \quad \forall I,$$

If  $n$  is such that  $L_{ij}$  does not depend on  $q_i^{[n]}$  for any  $I$ , then

$$\frac{\delta_{0i} L_{0i}}{\delta q_i^{[n]}} = 0 \quad \forall I \not\ni t_i,$$

$$\frac{\delta_{0j} L_{0j}}{\delta q_{It_j}^{[n]}} - \frac{\delta_{0i} L_{0i}}{\delta q_{It_i}^{[n]}} = 0 \quad \forall I.$$

# Contents

## ● Introduction

## ① Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries


## ② Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy

## ③ Summary

# Toda Lattice

Consider particles on a line: 1 discrete dimension, many continuous times



Denote  $q_1 = q_{t_1} = \frac{dq}{dt_1}$ ,  $q_{11} = q_{t_1 t_1} = \frac{d^2 q}{dt_1^2}$ , etc.

**Toda lattice:** exponential nearest-neighbour interaction

$$q_{11} = \exp(\bar{q} - q) - \exp(q - \underline{q}).$$

It is part of a hierarchy:

$$q_2 = q_1^2 + \exp(\bar{q} - q) + \exp(q - \underline{q})$$

$$q_3 = q_1^3 + (2q_1 + \underline{q}_1) \exp(q - \underline{q}) + (2q_1 + \bar{q}_1) \exp(\bar{q} - q)$$

$\dots$

# Toda lattice

Each member of the Toda hierarchy is Hamiltonian with Hamilton function of the form

$$H_i = \sum_{\alpha \in \mathbb{Z}} \mathcal{T}^\alpha h_i = \dots + \underline{h_i} + h_i + \overline{h_i} + \dots$$

Define  $L_{0j} = q_1 q_j - h_j$  and  $L_{ij}$  in such a way that  $\mathrm{d}\mathcal{L}$  will have a double zero on solutions of the hierarchy.

This is possible because the  $H_i$  are in involution.

We have a formula for  $L_{ij}$ , which this slide is too small to contain.

Semi-discrete Lagrangian 2-form  $\left( \sum_j L_{0j} \mathrm{d}t_j, \sum_{i,j} L_{ij} \mathrm{d}t_i \wedge \mathrm{d}t_j \right)$  with

$$L_{01} = \frac{1}{2} q_1^2 - \exp(\bar{q} - q)$$

$$L_{02} = q_1 q_2 - \frac{1}{3} q_1^3 - (q_1 + \bar{q}_1) \exp(\bar{q} - q)$$

$$L_{12} = -\frac{1}{4} (q_2 - q_{11} - q_1^2)^2$$

# Toda lattice

Euler-Lagrange equations:

$$\frac{\delta_{01} L_{01}}{\delta q} = 0 \quad \rightarrow \quad q_{11} = \exp(\bar{q} - q) - \exp(q - \underline{q})$$

$$\frac{\delta_{02} L_{02}}{\delta q_1} = 0 \quad \rightarrow \quad q_2 = q_1^2 + \exp(\bar{q} - q) + \exp(q - \underline{q})$$

$$\frac{\delta_{12} L_{12}}{\delta q} = 0 \quad \rightarrow \quad \frac{1}{2}q_{22} - q_{11}q_2 - 2q_{12}q_1 - \frac{1}{2}q_{1111} + 3q_1^2q_{11} = 0 \quad (*)$$

By construction, the variational principle is satisfied by the Toda hierarchy, so (\*) must be a consequence of the differential-difference equations

(\*) itself is an integrable PDE

The first two Toda equations can be written as

$$\exp(\bar{q} - q) = \frac{1}{2}(q_2 + q_{11} - q_1^2), \quad \exp(q - \underline{q}) = \frac{1}{2}(q_2 - q_{11} - q_1^2).$$

and form an auto-Bäcklund transformation for (\*)

## Toda lattice

At the next level we find

$$L_{03} = q_1 q_3 - \frac{1}{4} q_1^4 - a(q_1^2 + \bar{q}_1^2 + q_1 \bar{q}_1) - a\bar{a} + \frac{1}{2} a^2,$$

$$L_{13} = -a(\bar{q}_1^3 + 2a\bar{q}_1 + \bar{a}\bar{q}_1 + 2\bar{q}_1\bar{q}_{11} + q_1\bar{q}_{11} - \bar{q}_3 + aq_1 - \bar{a}q_1),$$

$$L_{23} = -a\left(\bar{q}_2(q_1^2 + \bar{q}_1^2 + q_1\bar{q}_1 + \underline{a} + a) + \bar{\bar{q}}_2\bar{a} + 2\bar{q}_1\bar{q}_{12} + q_1\bar{q}_{12} - \bar{q}_{13} \right. \\ \left. - \bar{q}_3(q_1 + \bar{q}_1) - q_1^2\bar{q}_1^2 - \underline{a}\bar{q}_1^2 + 2aq_1\bar{q}_1 - \bar{a}q_1^2 - a\bar{a} - \underline{a}\bar{a} - \underline{a}a - a^2\right)$$

where  $a = \exp(\bar{q} - q)$

Again we can use the multi-time Euler-Lagrange equations to obtain a PDE at a single lattice site:

$$\frac{\delta_{13} L_{13}}{\delta q} = 0 \quad \rightarrow \quad q_1^3 - 3q_1 q_{11} + 6q_1 a + q_{111} - q_3 = 0,$$

which can be simplified to

$$q_3 = -2q_1^3 + 3q_1 q_2 + q_{111}.$$

# Contents

## ● Introduction

### 1 Pluri-Lagrangian systems / Lagrangian multiforms

- Lagrangian 1-forms  $\rightarrow$  integrable ODEs
- Lagrangian 2-forms  $\rightarrow$  integrable PDEs
- Connections to Hamiltonian structures and variational symmetries

### 2 Differential-difference equations

- Semi-discrete Lagrangian multiforms
- The Toda Hierarchy

### 3 Summary

# Summary

- ▶ The **Lagrangian multiform** or **pluri-Lagrangian** principle describes symmetries and integrability.  
Applies to ODEs and PDEs, discrete and continuous.
- ▶ Closedness of the Lagrangian form, i.e.  $d\mathcal{L} = 0$ , is related to variational symmetries and Hamiltonians in involution.  
 $\delta d\mathcal{L} = 0$  is equivalent to the variational problem.
- ▶ We constructed **semi-discrete** Lagrangian 2-form for the Toda hierarchy. It reveals that integrable PDEs are hidden within.

# References

- Main ▶ Sleigh, V. [Semi-discrete Lagrangian 2-forms and the Toda hierarchy](#) [arXiv:2204.130633](#)
- Background ▶ Lobb, Nijhoff. [Lagrangian multiforms and multidimensional consistency](#). J. Phys. A. 2009.
- ▶ Yoo-Kong, Lobb, Nijhoff. [Discrete-time Calogero–Moser system and Lagrangian 1-form structure](#). J Phys. A. 2011.
  - ▶ Suris. [Variational formulation of commuting Hamiltonian flows: multi-time Lagrangian 1-forms](#). J. Geometric Mechanics, 2013
  - ▶ Suris, V. [On the Lagrangian structure of integrable hierarchies](#). In: [Advances in Discrete Differential Geometry](#), Springer. 2016.
  - ▶ Petrera, Suris. [Variational symmetries and pluri-Lagrangian systems in classical mechanics](#). J. Nonlin. Math. Phys., 2017.
  - ▶ V. [Hamiltonian structures for integrable hierarchies of Lagrangian PDEs](#) Open Communications in Nonlinear Mathematical Physics, 2021.

Thank you for your attention!